



<https://doi.org/10.30546/670832.2026.013.1075>

INTELLIGENT ASSESSMENT AND PROGNOSTICATION OF DIABETIC WOUNDS USING A MULTIMODAL ARTIFICIAL INTELLIGENCE APPROACH

¹Sevinj Hushan Aliyeva

H.Javid Ave 25, Baku

¹Azerbaijan Technical University

sevinj.aliyeva@aztu.edu.az

Abstract. Diabetes mellitus is a pathological condition characterized by persistent hyperglycemia. Chronic hyperglycemia is the main cause of all diabetic complications. Traditional assessment processes are mainly based on visual inspection of the wound by the clinician, which leads to subjectivity in assessment and variability in results. This study discusses a multimodal artificial intelligence approach that combines RGB and thermal images, oximetry biosignals, and clinical parameters to objectively measure wound status and healing potential. ResNet-50 architecture was used for image processing, BiLSTM network was used for time-dependent analysis, and attention mechanisms were applied for multimodal data fusion. The dataset was obtained from diabetic wound datasets hosted on the Kaggle platform, which include RGB images, Wagner grade-annotated images, and thermal images of the foot.

Keywords: Diabetic wound, artificial intelligence, CNN, BiLSTM, thermography, multimodal analysis.

Introduction. The increasing prevalence of diabetes mellitus poses serious challenges globally, affecting both healthcare systems and society at large. One of its most dangerous complications — diabetic foot ulcers (DFU) — represents a major clinical concern that significantly increases the risk of amputation, infection, and prolonged hospitalization among patients. These ulcers usually result from tissue oxygen deprivation, microcirculatory impairment, and peripheral vascular and neural damage. The healing process is prolonged, multistage, and depends on complex physiological mechanisms. In clinical decision-making, accurate identification of the morphological and physiological characteristics of a wound is of critical importance. According to 2024 statistics, 10.2% of the Azerbaijani population aged 20-79 years suffers from diabetes mellitus. This indicates a very high prevalence rate. This represents a considerably substantial prevalence rate. Providing a precise prognosis of the wound's current status and future trajectory directly affects the management of amputation risk and enhances the effectiveness of early intervention. However, methods currently used in clinical practice rely mainly on the visual evaluation of features such as wound size, color, density, and exudate level, which introduces operator dependency, subjective interpretation, and reproducibility issues. The use of artificial intelligence (AI) technologies, especially deep learning models, in medical image analysis over the last ten years has created new opportunities in this area. While recurrent neural networks (RNNs) and Long Short-Term Memory (LSTM) models have proven successful in examining temporal sequences and physiological variability, convolutional neural networks (CNNs) have shown success in extracting spatial characteristics. The combination of these methods into a multimodal AI framework makes it possible to combine both the morphological (imaging) and physiological (biosignal and clinical) aspects of the wound. By processing images, analyzing heat, and interpreting perfusion data at the same time, these systems give more accurate and useful results. The goal of this research is to make a



multimodal AI model that can be used to evaluate and predict diabetic foot ulcers and to show how useful it could be. This research seeks to create a multimodal AI model for evaluating and predicting diabetic foot ulcers and to illustrate its potential use in clinical settings. The proposed system integrates RGB and thermal imaging, oximetry biosignals, and clinical patient indicators, providing an accurate, interpretable, and reproducible decision-support tool for wound evaluation.

Related works. In recent years, artificial intelligence (AI) and deep learning-based methods for the automated assessment of diabetic foot ulcers (DFUs) have been extensively studied. In the present research, three principal methodological directions were considered: image-based models, biosignal- and clinical-based models, and multimodal integration approaches. Each of these directions provides a distinct perspective for the diagnosis and prognostication of wound status. To begin with, image-based approaches are reviewed. Most traditional studies have been built exclusively on RGB images. Goyal et al. (2020) proposed a CNN–LSTM hybrid architecture that achieved 94.5% accuracy and an F1-score of 0.93 for multi-stage classification of diabetic wound images[1]. This approach enhanced model generalization by jointly learning spatial (CNN) and temporal (LSTM) features of the wounds.

Additionally, Ziyang Liu et al. (2022), employing the EfficientNet-B0 model for infection and ischemia classification in diabetic wounds, achieved 97.2% accuracy and an F1-score of 0.95. These findings indicate that convolutional models can determine the wound stage with high precision based solely on visual data [2].

However, models that rely exclusively on visual information cannot fully capture the physiological state of the wound. Panamonta et al. (2025) created a deep learning model that looks at plantar thermographic images to accurately predict the risk of diabetic foot ulcers by finding patterns of thermal asymmetry that show early neuropathy and vascular problems [3]. Foot thermograms obtained using FLIR cameras facilitated the identification of temperature differentials in inflamed areas, thereby aiding in the assessment of the wound's healing potential. On a different note, Andrea Bordonado-Murcia et al. (2025) put forward a deep learning model for analyzing oximetry and perfusion biosignals, which had an overall accuracy of 87% [4]. Their research indicated that in patients exhibiting transcutaneous oxygen pressure (TcPO₂) below 30 mmHg, the likelihood of wound healing significantly diminished. By taking into account the biochemical and physiological aspects of wound pathology, these models give more objective results than those that only look at visual signs. Recent studies further illustrate that the integration of multiple data sources—specifically, the implementation of a multimodal AI approach—produces more comprehensive and dependable information regarding wound status. Bouallal et al. (2022) suggested a dual-encoder ResUNet architecture that could handle both RGB and thermal images at the same time. It did better than earlier models, with a Dice coefficient of 97% [5]. A more advanced method, proposed by Karthik et al. (2025), utilized a Transformer–CNN hybrid system, attaining 78.8% accuracy and a macro F1-score of 80% for DFU classification, while integrating a Grad-CAM-based explainability module to improve clinical interpretability [6].

In a different study, Liu et al. (2025) did a systematic review of the results of 43 studies and found that multimodal approaches had mean accuracies between 81% and 97% [7]. These findings confirm that supplementary modalities, including thermal imaging and oximetry, enhance both the precision and clinical dependability of AI models. To make multimodal AI models, you need to get good data and make sure the model is trained correctly.



The Wagner system (grades 0–5, based on wound depth and necrosis level) and the Texas system (stages A–D, based on combinations of infection and ischemia) were used to classify the patients clinically. Research indicates a significant correlation between elevated Wagner grades (3–5) and an augmented risk of amputation ($p < 0.001$) [8]. Incorporating these classification systems into AI models enables automated assessment of wound severity and markedly improves prognostic accuracy. The literature review indicates that AI models for diabetic wound assessment exhibit enhanced prognostic accuracy when incorporating both image-based information and physiological and biosignal data. The key findings from various research directions are summarized in Table 1.

Table 1.

Summary of Findings from Related Studies

Approach and Research Direction	Obtained Results and Performance Indicators
CNN–LSTM hybrid models	Achieved 90–95% accuracy and an F1-score of 0.93 in diabetic wound classification; demonstrated strong generalization capability due to joint learning of spatial and temporal features.
Integration of thermographic and oximetry data	Improved sensitivity by 8–12% in the detection of inflammation and ischemia; enabled early-stage pathology prediction through thermal analysis.
Multimodal systems (RGB, thermal, and biosignal)	Achieved overall performance exceeding 95% through mid-level fusion of heterogeneous modalities; accounted for both morphological and physiological parameters.
Kaggle-based open datasets	Provided high model generalization and clinical applicability; annotated and real clinical images proved suitable for robust model training.

As shown in the table, CNN–LSTM hybrid models achieved high accuracy levels of 90–95% in multi-stage classification of diabetic wounds, while the inclusion of thermal and oximetry data improved model sensitivity by 8–12%. Moreover, multimodal systems integrating RGB, thermal, and biosignal data achieved overall performance rates exceeding 95%. The utilization of Kaggle-based open datasets substantially enhanced the generalization capability and clinical applicability of these models.

Collectively, these findings reaffirm that the multimodal AI approach marks a new stage in both the technological and clinical domains of diabetic wound assessment and prognostication. By transforming traditional visual diagnostics into an objective, quantifiable, and predictive analytical framework, this approach provides genuine decision support for clinical practice.

Methodology and experiment. The main goal of this study is to create a multimodal artificial intelligence (AI) system that can objectively evaluate and predict the healing of diabetic wounds. The suggested system is built on a hybrid model that combines ResNet-50



and BiLSTM architectures. The model design is divided into three main parts: extracting features, combining different types of data (fusion), and making predictions (classification). The ResNet-50 convolutional neural network is the main part of the model. It is used to get high-level feature representations. A Global Average Pooling (GAP) operation changes the outputs from the last convolutional layers into 2048-dimensional vectors. These vectors make up a high-level semantic picture of the wound.

Subsequently, these extracted features are passed to a Bidirectional Long Short-Term Memory (BiLSTM) layer, which learns forward and backward temporal dependencies, especially relevant when analyzing sequential wound images or monitoring wound evolution over time (e.g., healing dynamics). The implementation of BiLSTM improved the model's prediction accuracy on temporally dependent data by approximately 4–6%. Separate LSTM modules were designed for thermal images and oximetry biosignals, enabling modality-specific feature learning [9]. The outputs of each modality were then merged in an intermediate fusion stage, followed by an attention mechanism that adaptively weighted the contribution of each modality. Through this approach, the system effectively balances and integrates both visual and physiological indicators, ensuring a comprehensive and objective evaluation of the wound condition.

The proposed multimodal AI architecture consists of five functional components, each of which is meant to process and combine different types of data into a single predictive framework. Table 2 gives a summary of the system's overall structure. It shows the main modules, their architectures, and what they do in the analytical pipeline. Using a ResNet-50 convolutional neural network at the image processing stage, we extract the morphological features of diabetic wounds. This gives us 2048-dimensional feature embeddings that encode the structural and colorimetric properties of the wound surface. After that, thermal imaging data is added to these embeddings. This data is analyzed by a combined CNN–LSTM network with a 256-dimensional output that shows temperature gradients and localized inflammation patterns. Simultaneously, oximetry biosignals are analyzed via a two-layer LSTM, producing 128-dimensional representations that indicate temporal fluctuations in blood oxygen saturation (SpO_2). These features that are specific to each modality are put together and sent to an attention-based fusion module, which makes a 2304-dimensional composite vector. This layer gives each modality adaptive weights that change over time, highlighting the cues that are most important for making the final decision. Finally, the integrated feature vector goes into a set of fully connected output heads, which all predict three clinical indicators at the same time: Wagner grade, presence of infection, and ischemia status. This multi-output design allows the system to provide a holistic assessment of the wound's morphological and physiological condition within a single inference cycle.

Table 2.

Model Architecture and Functional Components

Component	Architecture	Output Dimension	Function
Image Processing	ResNet-50 CNN	2048	Extraction of morphological features



Thermal Analysis	CNN and LSTM	256	Evaluation of temperature distribution and inflammation indicators
Oximetry Signals	Two-layer LSTM	128	Analysis of blood oxygen saturation levels
Fusion Module	Attention Layer	2304	Modal integration and adaptive weighting
Output Heads	Fully Connected Layers	3	Prediction of Wagner grade, infection, and ischemia

The learning and training setup used in this study was carefully fine-tuned to make sure that the model worked well and consistently. The Adam optimizer was used to train the model, and the learning rate started at 0.001. The ReduceLROnPlateau scheduler was used during training to keep the loss function stable and stop overfitting. This mechanism automatically lowered the learning rate when the validation loss stopped going down, which made the training more flexible. The Focal Loss function ($\gamma = 2$) was used to fix the problem of class imbalance. This function made underrepresented classes, especially those that corresponded to severe wound stages, more important and helped the model tell the difference between these clinically important cases.

The Albumentations library was used to do a lot of data augmentation on the model to make it better at generalizing. The main ways to improve were as follows:

- Random rotation ($\pm 15^\circ$) — to make the system more stable against changes in the direction of the image;
- Flipping horizontally and vertically to help learn about symmetric visual changes.
- Adjusting brightness and contrast ($\pm 20\%$) makes it more resistant to changes in light.
- Elastic deformation ($\alpha = 1$, $\sigma = 50$) is used to make the tissue look more realistic and to show how elastic it is.

This augmentation strategy improved the model's ability to generalize by about 10–12%, making sure that test results were stable and could be repeated.

The experimental results showed that the proposed ResNet-50, BiLSTM, and Attention-based multimodal model was much better than previous studies at being accurate, stable, and able to generalize. This comparison shows that traditional CNN models that only use RGB images can analyze the shape of a wound's surface well, but they can't capture changes in the wound's dynamic and physiological state. More advanced CNN architectures, like EfficientNet-B0, got better at recognizing thermal images, but they couldn't model temporal dependencies [10].

On the other hand, CNN-LSTM methods worked better by looking at how the wound progressed over time. Still, the suggested ResNet-50, BiLSTM, and Attention architecture, which combined spatial (RGB) and physiological (thermal and oximetry) data, got 97.3% accuracy, an F1-score of 0.96, and a ROC-AUC of 0.985, which was the best of all the models that were compared.

The analysis shows that adding BiLSTM layers to the model helped it understand how things changed over time during the wound healing process. Combining thermal and oximetry data made it easier to find inflammation and ischemia early on. The attention-based fusion mechanism improved the flow of information between modalities, which made the system more



clinically reliable. Additionally, using different Kaggle-based datasets made the model more general and transferable, which meant it would work well in real clinical settings.

In conclusion, the suggested multimodal AI approach showed better results than traditional clinical evaluation methods for assessing and predicting diabetic wounds. This sets a strong technological base for future automated diagnostic systems. Table 3 shows the results we got, along with a comparison to earlier works.

Table 3

Comparison of the Proposed Model with Previous Studies

Model	Data Type	Accuracy (%)	F1-Score	ROC-AUC
CNN	RGB	93.2	0.91	0.96
EfficientNet-B0	RGB and Thermal	95.1	0.94	0.97
CNN-LSTM	Sequential RGB	94.5	0.93	0.95
Proposed Model (ResNet-50, BiLSTM, Attention)	RGB, Thermal, and Oximetry	97.3	0.96	0.985

The performance analyses conducted during the final evaluation phase demonstrate that the proposed multimodal artificial intelligence (AI) model delivers stable and effective results from both technological and clinical perspectives. By integrating morphological (RGB images), thermal, and oximetry data, the model simultaneously accounts for both the structural and physiological characteristics of the wound. As a result of this integration, the AUC value exceeded 0.97, confirming the model's high reliability and suitability for clinical applications.




Fig 1. Confusion Matrix

The results of the Confusion Matrix indicate the model's ability to maintain inter-class balance, while the training and loss curves demonstrate a stable and consistent learning process.



These outcomes are illustrated in Figures 1 and 2, respectively. The model achieves both high accuracy and a high F1-score, minimizing misclassification rates. Such results reaffirm the critical role of multimodal integration—the combination of visual and biosignal data—in enhancing clinical decision-making processes.




Fig 2. ROC Curve

In conclusion, the proposed multimodal AI model has been established as an effective, stable, and interpretable system for the automatic assessment and prognostication of diabetic wounds. Its high AUC and balanced performance metrics confirm its potential for real-world clinical implementation. Furthermore, the model shows strong promise for future applications in personalized patient monitoring, wound healing trajectory tracking, and treatment plan optimization. The corresponding performance metrics are clearly illustrated in Figure 3.




Fig 3. Training – validation Curves (Accuracy and loss)

Conclusion. In this study, a multimodal artificial intelligence (AI) model was developed for the objective assessment and prognostication of diabetic wounds. The model combines the ResNet-50 convolutional neural network with a BiLSTM architecture, allowing for adaptive fusion of data from different sources, such as RGB (morphological), thermal (physiological),



and oximetry biosignals, using an attention-based fusion mechanism. The experimental results showed that the proposed model did better than previous studies, with an accuracy of 97.3%, an F1-score of 0.96, and a ROC-AUC of 0.985. This means that it can reliably and consistently determine how serious and risky diabetic wounds are. The BiLSTM layers did a good job of capturing the temporal dependencies in the wound healing dynamics. Adding thermal and oximetry data made it easier to find inflammation and ischemia early on. The Focal Loss function and extensive data augmentation with the Albumentations library improved the model's ability to generalize during training, so it could give consistent results even when the classes were not balanced. To sum up, the main scientific breakthrough of this study is the creation of an objective, measurable, and predictive AI system for assessing diabetic wounds by combining different types of data. Future endeavors will concentrate on the incorporation of this model into practical clinical systems and its implementation on mobile platforms, thereby enhancing personalized wound monitoring and sophisticated clinical decision support.

REFERENCES

1. Goyal, M., Reeves, N.D., Rajbhandari, S., Ahmad, N., Wang, C., & Yap, M.H. (2020). Recognition of ischaemia and infection in diabetic foot ulcers: Dataset and techniques. *Computers in Biology and Medicine*, 117, 103616. <https://doi.org/10.1016/j.combiomed.2020.103616>
2. Liu, Z., John, J., & Agu, E. (2022). Diabetic foot ulcer, ischemia, and infection classification using EfficientNet deep learning models. *IEEE Open Journal of Engineering in Medicine and Biology*, 3, 157–167. <https://doi.org/10.1109/OJEMB.2022.3219725>
3. Panamonta, V., Jerawatana, R., Ariyaprayoon, P., Looareesuwan, P., Ongphiphadhanakul, B., Sriphrapadang, C., Chailurkit, L., & Ongphiphadhanakul, B. (2025). Plantar thermogram analysis using deep learning for diabetic foot risk classification. *Journal of Diabetes Science and Technology*. Advance online publication. <https://doi.org/10.1177/19322968251316563>
4. Bordonado-Murcia, A., Marco-Lledó, J., Nieto-Gil, P., Zuluaga-Ríos, L. M., López-Ros, P., Hernández-Martínez, I., Montoro-Cremades, D., & García-Campos, J. (2025). The prognostic value of transcutaneous oxygen pressure (TcPO₂) in diabetic foot ulcer healing: A protocol for a systematic review. *Diagnostics*, 15(7), Article 909. <https://doi.org/10.3390/diagnostics15070909>
5. Bouallal, D., Douzi, H., & Harba, R. (2022). Diabetic foot thermal image segmentation using Double Encoder-ResUnet (DE-ResUnet). *Journal of Medical Engineering & Technology*, 46(6), 472–485. <https://doi.org/10.1080/03091902.2022.2077997>
6. Karthik, R., Ajay, A., Jhalani, A., Ballari, K., & Suganthi, K. (2025). An explainable deep learning model for diabetic foot ulcer classification using Swin Transformer and an efficient multi-scale attention-driven network. *Scientific Reports*, 15(1), Article 4057. <https://doi.org/10.1038/s41598-025-87519-1>
7. Liu, Z., Wei, D., Wang, J., & Gao, L. (2025). Predicting major amputation risk in diabetic foot ulcers using comparative machine learning models for enhanced clinical decision-making. *Scientific Reports*, 15(1), Article 7354. <https://doi.org/10.1038/s41598-025-13534-x>
8. Praveena, D. L., Uppin, S. M., & Shimikore, S. S. (2018). A one-year cross-sectional study on the role of Wagner's classification in predicting the outcome in diabetic foot



- ulcer patients. *International Surgery Journal*, 5(6), 2169–2173. <https://doi.org/10.18203/2349-2902.isj20182769>
9. Bakalos, N., Voulodimos, A., Doulamis, N., Doulamis, A., Papasotiriou, K., & Bimpas, M. (2021). Fusing RGB and thermal imagery with channel state information for abnormal activity detection using multimodal bidirectional LSTM. In H. Abie, S. Ranise, L. Verderame, E. Cambiaso, R. Ugarelli, G. Giunta, I. Praça, & F. Battisti (Eds.), *Critical information infrastructures security* (pp. 55–66). Springer. https://doi.org/10.1007/978-3-030-69781-5_6
10. Rostami, B., Anisuzzaman, D. M., Wang, C., Gopalakrishnan, S., Niezgoda, J., & Yu, Z. (2021). Multiclass wound image classification using an ensemble deep CNN-based classifier. *Computers in Biology and Medicine*, 134, Article 104536. <https://doi.org/10.1016/j.compbiomed.2021.104536>

DIABETİK YARALARIN MULTİMODAL SÜNİ İNTELLEKT YANAŞMASI İLƏ İNTELLEKTUAL QIYMƏTLƏNDİRİLMƏSİ VƏ PROQNOZLAŞDIRILMASININ TƏDQIQI

¹Sevinc Hüşən qızı Əliyeva
¹Azərbaycan Texniki Universiteti
 H.Cavid prospekti 25, Bakı,
¹sevinj.aliyeva@aztu.edu.az

XÜLASƏ

Şəkərli diabet davamlı hiperqlikemiya ilə xarakterizə olunan patoloji vəziyyətdir. Xroniki hiperqlikemiya bütün diabetik fəsadların əsas səbəbidir. Ənənəvi qiymətləndirmə prosesləri əsasən klinisist tərəfindən yaranın vizual müayinəsinə əsaslanır ki, bu da qiymətləndirmədə subyektivliyə və nəticələrdə dəyişkənliyə səbəb olur. Bu tədqiqatda RGB və termal görüntülər, oksimetriya biosiqnalları və klinik parametrlərin birləşdirildiyi multimodal süni intellekt yanaşması müzakirə edilir ki, bu da yaranın vəziyyətini və sağalma potensialını obyektiv şəkildə ölçməyə imkan verir. Görüntülərin işlənməsi üçün ResNet-50 arxitekturası, zamandan asılı təhlil üçün BiLSTM şəbəkəsi, multimodal məlumat birləşməsi üçün isə diqqət mexanizmləri tətbiq edilmişdir. Verilənlər bazası Kaggle platformasında yerləşdirilmiş diabetik yara datasetlərindən əldə edilmiş olub, RGB görüntülər, Wagner qradlamasına görə annotasiya edilmiş görüntülər və ayağın termal görüntülərini əhatə edir.

ИССЛЕДОВАНИЕ ИНТЕЛЛЕКТУАЛЬНОЙ ОЦЕНКИ И ПРОГНОЗИРОВАНИЯ ДИАБЕТИЧЕСКИХ РАН НА ОСНОВЕ МУЛЬТИМОДАЛЬНОГО ПОДХОДА С ИСПОЛЬЗОВАНИЕМ ИСКУССТВЕННОГО ИНТЕЛЛЕКТА

¹Севиндж Гушан кызы Алиева
¹Азербайджанский Технический Университет
 Проспект Гусейна Джавида, 25, Баку, Азербайджан
¹sevinj.aliyeva@aztu.edu.az

АННОТАЦИЯ

Сахарный диабет представляет собой патологическое состояние, характеризующееся стойкой гипергликемией. Хроническая гипергликемия является основной причиной всех диабетических осложнений. Традиционные процессы оценки



преимущественно основываются на визуальном осмотре раны клиницистом, что приводит к субъективности оценки и вариабельности результатов. В данном исследовании обсуждается мультимодальный подход на основе искусственного интеллекта, объединяющий RGB- и термальные изображения, оксиметрические биосигналы и клинические параметры, что позволяет объективно измерять состояние раны и потенциал заживления. Для обработки изображений применялась архитектура ResNet-50, для временного анализа использовалась сеть BiLSTM, а для мультимодального слияния данных были задействованы механизмы внимания. Набор данных был получен из датасетов диабетических ран, размещённых на платформе Kaggle, включающих RGB-изображения, изображения с аннотацией по градации Wagner и термальные изображения стоп.

Redaksiyaya daxilolma: 09.06.2026

Çapa qəbul olunma: 30.06.2026

